

## Fixed Point Iterations for Multi-Valued Mapping in Uniformly Smooth Banach Space

### Abstract

Let  $K$  be nonempty closed convex bounded subset of a real uniformly smooth Banach space with modulus of smoothness of power type  $q > 1$  and  $T: K \rightarrow P(K)$  a multi-valued quasi-contractive mapping has a fixed point. We proved that the sequences of Mann and Ishikawa iterate converges to a fixed point of  $T$ . The result generalizes and extends former result proved by Sastry and Babu [17].

**Keywords:** Multi-value mapping, Mann iterates, Ishikawa iterates, fixed points, uniformly smooth Banach space.

### Introduction

In 1974, Ćirić [8] introduced a new mapping which is known as quasi-contractive mapping. Let  $K$  be a nonempty subset of normed linear space  $X$  and  $T$  is a mapping of  $K$  in to itself. Then  $T$  is said to quasi-contractive ([8]), if there exists a constant  $K \in [0, 1)$  such that

$$\|Tx - Ty\| \leq k \max\{\|x - y\|, \|x - Tx\|, \|y - Ty\|, \|x - Ty\|, \|y - Tx\|\} \quad (1.1)$$

for all  $x, y \in K$ . In [16], Rhoades showed that the contractive definition (1.1) apart from being an obvious generalization of the well known contractive mapping is one of the most general contractive definitions for which Picard iteration give a unique fixed point.

In [15], Rhoades examined the following two iteration processes:

(a) Mann iteration process defined as follows:

For  $K$  a convex subset of Banach space  $X$  and  $T$  be a nonlinear mapping of  $K$  into itself, the sequence  $\{x_n\}$  in  $K$  defined by

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n Tx_n,$$

Where  $\{\alpha_n\}$  is sequence in  $[0, 1]$  satisfy certain restriction.

(b) Ishikawa iteration process defined as follows:

With  $K$  and  $T$  as in (a), the sequence  $\{x_n\}$  in  $K$  defined by

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n Ty_n, \\ y_n = (1 - \beta_n)x_n + \beta_n Tx_n,$$

Where  $\{\alpha_n\}$  and  $\{\beta_n\}$  are sequences in  $[0, 1]$  satisfying certain restriction.

<sup>o</sup>2000 AMS Subject Classifications: 47H10, 54H25.

The above iteration processes have been studied by several authors (see e.g., Chidume ([3, 4, 5, 6, 7]), Ishikawa [9], Mann [11], Qihou ([14]), Rhoades ([15]), for approximating solutions of several nonlinear operator equations. Moreover, it is well known that the two schemes may exhibit different behaviours for different classes of nonlinear mappings (see e.g., Rhoades [15]).

The study of fixed point problems for multi-valued mappings was initiated by Kakutani in 1941. Later several authors have worked on this setting. For instance, Nadler jr. [13] has generalized the Banach contraction principle theorem to multi-valued mappings. Subsequently Assad and Kirk [1] have worked with contractive type mapping and Markin [12] pursued the theory to nonexpansive mappings in multi-valued setting and many others in these directions.

Let  $X$  is Banach space. A subset of  $K$  is called proximal if for each  $x \in X$ ; there exists an element  $k \in K$  such that

$$d(x, k) = d(x, k) = \inf \{\|x - y\| : y \in K\},$$

It is well known that every closed convex subset of a uniformly convex Banach space is proximal. We denote the family of all nonempty bounded proximal subset of  $K$  by  $P(K)$ . Let  $A, B$  be two bounded subsets of  $X$ . The Hausdorff distance between  $A$  and  $B$  defined by

$$H(A, B) = \max\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A) : A, B \in P(K) \},$$

**Samir Dashputre**

Desig.....

Deptt. of Applied Mathematics,  
Shri Shankaracharya Groups of  
Institutions,  
Junwani, Bhilai, (C.G)

**Yogesh kumar Sahu**

Desig.....

Deptt. of Applied Mathematics,  
Shri Shankaracharya Groups of  
Institutions,  
Junwani, Bhilai, (C.G)

**(Please fill above detail)**

Where  $d(a, b) = \inf \{ \|a - b\| : b \in B \}$  is the distance from the point a to the set B.

Recently, Sastry and Babu [17] extend the convergence results from single valued maps to multi-valued maps by defining the analogs of Mann and Ishikawa iterates for multi-valued maps with a fixed point.

**Definition 1.1.**

(Sastry and Babu [17]). Let K be nonempty convex subset of X,

$T: K \rightarrow P(K)$  a multi-valued mapping and  $p \in F(T)$ .

(A) The sequence of Mann iterates is defined by  $\{x_n\}$  in K,

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n u_n, \alpha_n \in [0,1]; \forall n \geq 0$$

Where  $u_n \in Tx_n$  is such that  $\|u_n - p\| = d(p, Tx_n)$ .

(B) The sequence of Ishikawa iterates is defined by  $\{x_n\}$  in K,

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n z_n, \alpha_n \in [0,1]; \forall n \geq 0$$

Where  $z_n \in Ty_n$  is such that  $\|z_n - p\| = d(p, Ty_n)$ .

$$y_n = (1 - \beta_n)x_n + \beta_n z_n, \beta_n \in [0,1]; \forall n \geq 0$$

Where  $z_n \in Tx_n$  is such that  $\|z_n - p\| = d(p, Tx_n)$ .

Fixed Point Iterations for Multi-valued Mapping in Uniformly Smooth Banach Space3

**Definition 1.2**

(Sastry and Babu [17]). A multi-valued  $T: K \rightarrow P(K)$  is said to be quasi-contractive if for some constant k,  $0 \leq k < 1$

$$\|Hx - Hy\| \leq k \max \{ \|x - y\|, d(x, Tx), d(y, Ty), d(y, Tx); d(x, Ty) \} \quad (1.2)$$

For all  $x, y \in C$ . A point x is called fixed point of T if  $x \in Tx$ .

In [17] Sastry and Babu proved the following theorem:

**Theorem 1.3**

(Sastry and Babu [17], Theorem 9, p.824).

Let K be a closed convex bounded subset of Hilbert space X. Suppose  $T: K \rightarrow P(K)$  is quasi contractive (1.2) and has a fixed point  $p \in F(T)$ . Let  $\{\alpha_n\}$  and  $\{\beta_n\}$  be real sequences such that  $0 \leq \alpha_n, \beta_n < 1$  for all n and  $\beta_n \rightarrow 0$  as  $n \rightarrow \infty$  with  $\delta \leq \alpha_n \leq 1 - k^2$  for some  $\delta > 0$ . Then sequence of Ishikawa iterates defined by (B) converges to p.

The main purpose of this paper extend the convergence result of Sastry and Babu [17] from Hilbert space to uniformly smooth Banach space with modulus of smoothness of power type  $q > 1$ , and also our result extends the convergence result of Chidume and Osilike [7] from single-valued maps to multi-valued maps.

**Preliminaries**

We begin with the following definition and lemmas:

Let X be a Banach space. The modulus of smoothness of X is the function

$$\rho_X : [0, \infty) \rightarrow [0, \infty)$$

Defined by

$$\rho_X(\tau) := \sup \{ \frac{1}{2} (\|x + y\| + \|x - y\|) - 1 : \|x\| \leq 1; \|y\| \leq \tau \}.$$

X is said to be uniformly smooth if:

$$\lim_{\tau \rightarrow 0} \frac{\rho_X(\tau)}{\tau} = 0:$$

A uniformly smooth Banach space X is said to have smoothness of power type  $q > 1$  if there exists a constant  $c > 0$  such that  $\rho_X(\tau) \leq c\tau^q$ .

Let  $q > 1$ ,  $J_q$  denotes the generalized duality mapping from X to  $2^{X^*}$  given by

$$J_q(x) = \{ f^* \in X^* : \langle x, f^* \rangle = \|x\|^q, \|f^*\| = \|x\|^{q-1} \},$$

Where  $X^*$  denotes the dual space of X and  $\langle \cdot, \cdot \rangle$  denote the generalized duality pairing. In particular  $J = J_2$  is called the *normalized duality mapping* and  $J_q = \|x\|^{q-2} J(X)$  if  $x \neq 0$ . It is well known that (see e.g., Xu [19]) X is uniformly smooth, if and only if J (and hence  $J_q$ ) is single valued and uniformly continuous on any bounded subset of X.

In this sequel, we shall need the following lemmas.

**Lemma 2.1**

([18], Corollary 1, p. 1130) Let X be a uniformly smooth Banach space. Then X has modulus of smoothness of power type  $q > 1$  if and only if there exists a constant  $c > 0$  such that

$$\|x + y\|^q \leq \|x\|^q + q \langle y, J_q(x) \rangle + c \|y\|^q \quad (2.1)$$

for all  $x, y \in X$ .

For Hilbert space  $q = 2$ ,  $c = 1$  and equality holds. For  $p \geq 2, L_p$  or  $(l_p)$  spaces have modulus of smoothness of power type  $q = 2$  and (2.1) is satisfied with  $c = (p - 1)$  (see e.g., Xu [18]).

**Lemma 2.2**

(Chidume and Osilike [7], p.206) Let X be a uniformly smooth Banach space with modulus of smoothness of power type  $q > 1$ . Then for all  $x, y, z \in X$  and  $t \in [0, 1]$ ,

$$\|tx + (1-t)y - z\|^q \leq [1-t(q-1)]\|y-z\|^q + tc\|x-z\|^q - t[1-t^{q-1}c]\|x-y\|^q, \quad (2.2)$$

Where  $c > 0$  is a constant appearing in (2.1).

**Lemma 2.3**

(Liu [10], p.118) Let  $\{a_n\} n \geq 0, \{b_n\} n \geq 0, \{c_n\} n \geq 0$ , and  $\{t_n\} n \geq 0$ , be four sequences of nonnegative numbers satisfying the inequality:

$$a_{n+1} \leq (1 - t_n)a_n + t_n b_n + c_n \text{ for all } n \geq 0,$$

where  $\{t_n\} n \geq 0 \subset [0,1], c_n < \infty$  and  $\lim_{n \rightarrow \infty} b_n \sum_{n=0}^{\infty} t_n = \infty, \sum_{n=0}^{\infty} c_n = 0$ . Then

$$\lim_{n \rightarrow \infty} a_n = 0.$$

**Main Results**

**Theorem 3.1.**

Let X be a real uniformly smooth Banach space with modulus of smoothness of power type  $q > 1$ . Let K be a nonempty, closed convex and bounded subset of X. Suppose  $T: K \rightarrow P(K)$  is a multi-valued quasi-contractive mapping and has fixed point p. Let

$$(i) \quad c \geq 1, q - 1 \geq ck^q,$$

where c is the constant appearing in (2.1).

Let  $\{\alpha_n\}$  and  $\{\beta_n\}$  real sequences in  $[0,1]$  satisfying

the condition:

- (ii)  $\sum_{n=1}^{\infty} \alpha_n = \infty$ ,
- (iii)  $\lim_{n \rightarrow \infty} \beta_n = 0$
- (iv)  $c\alpha_n^{q-1} - ck^q\beta_n(q-1) \leq 1 - ck^q$ ,
- (v)  $c\beta_n^{q-1} \leq \frac{2-ck^q}{2}$ .

Then for any  $x_0 \in K$ , the sequence  $\{x_n\}$  defined by (B) converges to a fixed point of T.

**Proof**

Let p be fixed point of T then by using Lemma 2.2 and (B), we have

Fixed Point Iterations for Multi-valued Mapping in Uniformly Smooth Banach Space5

$$\begin{aligned} \|x_{n+1} - p\|^q &= \|(1-\alpha_n)x_n + \alpha_n z'_n - p\|^q \\ &\leq (1-\alpha_n)(q-1)\|x_n - p\|^q + c\alpha_n \|z'_n - p\|^q \\ &\quad - \alpha_n(1-\alpha_n^{q-1})\|x_n - z'_n\|^q. \end{aligned} \quad (3.2)$$

Again

$$\begin{aligned} \|z'_n - p\| &= d(p, Ty_n) \\ &\leq \max_{z \in T_p} d(z, Ty_n) \\ &\leq H(T_p, Ty_n). \end{aligned}$$

Therefore,

$$\|z'_n - p\|^q = H^q(T_p, Ty_n) \leq k^q \max\{\|y_n - p\|^q, d^q(y_n, Ty_n), d^q(p, Ty_n)\} \quad (3.2)$$

If  $d^q(p, Ty_n)$  is maximum, then

$$\begin{aligned} H^q(T_p, Ty_n) &\leq k^q d^q(p, Ty_n) \\ &\leq k^q \max_{z \in T_p} d^q(z, Ty_n) \\ &\leq k^q H^q(T_p, Ty_n), \end{aligned}$$

so that  $0 \leq \|z'_n - p\|^q \leq H^q(T_p, Ty_n) = 0$ . Hence, from (3.2), we get always

$$\begin{aligned} \|z'_n - p\| &\leq K^q \max\{\|y_n - p\|^q, d^q(y_n, Ty_n)\}, \\ &\leq k^q [\|y_n - p\|^q + d^q(y_n, Ty_n)] \end{aligned} \quad (3.3)$$

Using Lemma 2.2, (3.3) and (B), we have

$$\begin{aligned} \|y_n - p\|^q &= \|(1-\beta_n)x_n + \beta_n z_n - p\|^q \\ &\leq (1-\beta_n)(q-1)\|x_n - p\|^q + \beta_n c \|z_n - p\|^q \\ &\quad - \beta_n(1-\beta_n^{q-1}c)\|x_n - z_n\|^q \end{aligned} \quad (3.4)$$

$$\begin{aligned} d^q(y_n, Ty_n) &\leq \|y_n - z'_n\|^q \\ &\leq \|(1-\beta_n)x_n + \beta_n z_n - z'_n\|^q \\ &\leq (1-\beta_n)(q-1)\|x_n - z'_n\|^q + \beta_n c \|z_n - z'_n\|^q \\ &\quad - \beta_n(1-\beta_n^{q-1}c)\|x_n - z_n\|^q. \end{aligned} \quad (3.5)$$

Using (3.3),(3.4) and (3.5), we have

$$\begin{aligned} \|z'_n - p\|^q &\leq k^q (1-\beta_n)(q-1)\|x_n - p\|^q \\ &\quad + k^q \beta_n c \|z_n - p\|^q \\ &\quad + k^q \beta_n c \|z_n - z'_n\|^q k^q (1-\beta_n)(q-1)\|x_n - z'_n\|^q \\ &\quad - 2k^q \beta_n (1-\beta_n^{q-1}c)\|x_n - z_n\|^q. \end{aligned} \quad (3.6)$$

Similar to the inequality (3.3), we get that

$$\begin{aligned} \|z_n - p\|^q &= d^q(p, Tx_n) \\ &\leq H^q(T_p, Tx_n) \\ &\leq k^q [\|x_n - p\|^q + d^q(x_n, Tx_n)]. \end{aligned} \quad (3.7)$$

From (3.6) and (3.7), we get

$$\begin{aligned} \|z'_n - p\|^q &\leq k^q \|x_n - p\|^q + k^q \beta_n c \|z_n - z'_n\|^q \\ &\quad + k^q (1-\beta_n)(q-1)\|x_n - z'_n\|^q \\ &\quad - k^q \beta_n (2-2\beta_n^{q-1}c-ck^q)d^q(x_n, Tx_n). \end{aligned} \quad (3.8)$$

From (3.1) and (3.8), we have

$$\begin{aligned} \|x_{n+1} - p\|^q &= (1-\alpha_n)(q-1)\|x_n - p\|^q - \alpha_n(1-\alpha_n^{q-1}c)\|x_n - z'_n\|^q \\ &\quad + c\alpha_n [k^q \|x_n - p\|^q + k^q \beta_n c \|z_n - z'_n\|^q \\ &\quad + k^q (1-\beta_n)(q-1)\|x_n - z'_n\|^q \\ &\quad - k^q \beta_n (2-2\beta_n^{q-1}c-ck^q)d^q(x_n, Tx_n)] \\ &\leq (1-\alpha_n)(q-1-ck^q)\|x_n - p\|^q + c^2 \alpha_n \beta_n k^q \|z_n - z'_n\|^q \\ &\quad - ck^q \alpha_n \beta_n (2-2\beta_n^{q-1}c-ck^q)d^q(x_n, Tx_n) \\ &\quad - \alpha_n [1-\alpha_n^{q-1}ck^q(1-\beta_n)(q-1)]\|x_n - z'_n\|^q. \end{aligned} \quad (3.9)$$

As  $\beta_n \rightarrow 0$  as  $n \rightarrow \infty$ , there exists a positive integer  $N_1$  such that

$$c\beta_n^{q-1} \leq \frac{2-ck^q}{2} \quad \forall n \geq N_1$$

so that

$$2-2\beta_n^{q-1}c-ck^q \geq 0 \quad \forall n \geq N_1.$$

Fixed Point Iterations for Multi-valued Mapping in Uniformly Smooth Banach Space7

Also we have

$$\alpha_n^{q-1}c - ck^q\beta_n(q-1) \leq 1 - ck^q,$$

Implies that

$$1 - \alpha_n^{q-1}c - ck^q(1-\beta_n)(q-1) \geq 0,$$

For all  $n \geq N_1$ . Consequently, from (3.9), we get that for sufficiently large n

$$\begin{aligned} \|x_{n+1} - p\|^q &\leq (1-\alpha_n)(q-1-ck^q)\|x_n - p\|^q \\ &\quad + \alpha_n \beta_n c^2 k^q D, \end{aligned} \quad (3.10)$$

Where D is the diameter of C. Now by using lemma 2.3, the sequence  $\{x_n\}$  converges to p as  $n \rightarrow \infty$ .

**Remark 3.2**

For Hilbert spaces  $q = 2$  and  $c = 1$ , so that if we set  $q = 2$  and  $c = 1$  in Theorem 3.1, then condition  $(q-1-ck^q)$  reduces to  $(1-k^2) < 1$ . Moreover, conditions (iii), (iv), and (v) reduce to exactly the same condition of theorem 9 of Sastry and Babu [17].

**Theorem 3.3**

Let X be a real uniformly smooth Banach space with modulus of smoothness of power type  $q > 1$ . Let k be a closed convex and bounded subset of X. Suppose  $T : K \rightarrow P(K)$  is a quasi-contractive and has fixed point p. Let  $\{\alpha_n\}$  be a real sequence satisfying:

- (i)  $0 \leq \alpha_n < 1$  for all  $n \geq 0$
- (ii)  $\lim_{n \rightarrow \infty} \alpha_n = 0$ ,
- (iii)  $\sum_{n=0}^{\infty} \alpha_n = \infty$ ,

Then the sequence  $\{x_n\}$  defined by (A), converges to a fixed point of T.

**Proof.** By using Lemma 2.2 and (A), we get

$$\begin{aligned} \|x_{n+1} - p\|^q &= \|(1-\alpha_n)x_n + \alpha_n u_n - p\|^q \\ &\leq (1-\alpha_n)(q-1)\|x_n - p\|^q + \alpha_n c \|u_n - p\|^q \\ &\quad - \alpha_n(1-\alpha_n^{q-1}c)\|x_n - u_n\|^q \end{aligned} \quad (3.11)$$

Similar to inequality, we get

$$\begin{aligned} \|u_n - p\|^q &= d^q(p, Tx_n) \\ &\leq H^q(T_p, Tx_n) \\ &\leq k^q \|x_n - p\|^q + d^q(x_n, Tx_n). \end{aligned} \quad (3.12)$$

Then From (3.11) and (3.12), we get

$$\begin{aligned} \|x_{n+1} - p\|^q &= (1-\alpha_n)(q-1)\|x_n - p\|^q + \alpha_n c k^q \|x_n - p\|^q \\ &\quad + c\alpha_n k^q d^q(x_n, Tx_n) - \alpha_n(1-\alpha_n^{q-1}c)\|u_n - x_n\|^q \\ &\leq (1-\alpha_n)(q-1-ck^q)\|x_n - p\|^q \\ &\quad - \alpha_n(1-c\alpha_n^{q-1}-ck^q)d^q(x_n, Tx_n). \end{aligned} \quad (3.13)$$

Condition (ii) implies for some  $N_0$  sufficient large  $N_0 \in N$ ,  $(1 - \alpha_n(q - 1 - ck^q)) < 1$  and  $(1 - \alpha_n(q - 1 - ck^q)) \geq 0$  so that from (3.13), we have

$$\|x_{n+1} - p\|^q \leq (1 - \alpha_n(q - 1 - ck^q)) \|x_n - p\|^q. \quad (3.14)$$

Iteration of inequality (3.14) from  $j = N_0$  to  $N$  yields,

$$\|x_{N+1} - p\|^q \leq \prod_{j=N_0}^N (1 - \alpha_j(q - 1 - ck^q)) \|x_{N_0} - p\|^q = 0$$

as  $n \rightarrow \infty$  by condition (iii). Hence  $x_n \rightarrow p$  as  $n \rightarrow \infty$ .

#### Conclusion

(4.1) We conclude that, by theorems 3.1 and 3.3, either Mann or Ishikawa iterates can be used to approximate the fixed point for multi-valued quasi-contractive mapping in real uniformly smooth Banach space with modulus of smoothness of power type  $q > 1$ .

(4.2) Theorems 3.1 and 3.3 extends Theorem [9] of Sastry and Babu [17] from Hilbert space to more general Banach space.

(4.3) Theorems 3.1 and 3.3 extends Theorems 1 and 2 of Chidume and Osilike [7] from single valued maps to multi-valued maps. Fixed Point Iterations for Multi-valued Mapping in Uniformly Smooth Banach Space 9.

#### References

1. N.A. Assad and W.A. Kirk, Fixed point theorems for set valued mapping of contractive type, Pacific J. Math. 43 (1972), 553–562.
2. F.E. Browder, The fixed point theory of multi-valued mapping in topological vector spaces, Math Ann. 177 (1968), 283–301.
3. C.E. Chidume, on the Ishikawa fixed point iteration for quasi-contractive mappings, J. Nigerian Math. Soc. 4 (1985), 1–11.
4. C.E. Chidume, on the Ishikawa and Mann iteration methods for quasi-contractive mappings, J. Nigerian Math. Soc. 7 (1988), 1–9.
5. C.E. Chidume, Fixed point iterations for certain classes of nonlinear mappings, Applicable Anal. 27(1–3) (1988), 31–45.
6. C.E. Chidume, Approximation of fixed points of quasi-contractive mappings in  $L_p$  Spaces, Indian J. Pure. Appl. Math. 22(4) (1991), 273–286.
7. C.E. Chidume and M. O. Osilike, Fixed point iterations for quasi-contractive maps in uniformly smooth Banach spaces, Bull Korean Math. Soc. 30 (2) (1993), 201–212.
8. L.B. Ćirić, A generalization of Banach's contraction principle, Proc. Amer. Math. Soc. 45(2) (1974), 267–273.
9. S. Ishikawa, Fixed points by a new iteration method, Proc. Amer. Math. Soc. 44(1) (1974), 147–150.
10. L.S. Liu, Ishikawa and Mann iterative process with errors for nonlinear strongly accretive mappings in Banach spaces, J. Math Anal. Appl. 194 (1995), 114–125.
11. W.R. Mann, Mean value methods in iteration, Proc. Amer. Math. Soc. 4(1953), 506–510.
12. J.T. Markin, A Fixed point theorem for set-valued mapping, Bull. Amer. Math. Soc. 74 (1968), 639–640.
13. J.R. S.B. Nadler, Multi-valued contraction

mappings, Pacific J. Math. (1969), 457–588.

14. L. Qihou, A convergence theorem of the sequence of Ishikawa iterates for quasi-contractive mappings, J. Math. Anal. Appl. 146 (1990), 301–305.
15. B.E. Rhoades, comments on two fixed point iteration methods, J. Math. Anal. Appl. 56 (2) (1976), 741–750.
16. B.E. Rhoades, A comparison of various definitions of contractive mappings, Trans. Amer. Math. Soc. 226 (1977), 257–290.
17. K.P.R. Sastry and G.V.R. Babu, convergence of Ishikawa iterates for multi-valued mapping with fixed point, Czechoslovak Math. J. 55 (2005), 817–826.
18. H.K. Xu, Inequalities in Banach spaces with applications, Nonlinear Anal. 16(2) (1991), 1127–1138.
19. Z.B. Xu and C.F. Roach, Characteristic inequalities of uniformly convex and uniformly smooth Banach spaces, J. Math. Anal. Appl., 157 (1991), 189–210.