

Medicinal Weeds of Rabi (Winter) Crops of Tehsil Sunderbani District Rajouri (Jammu & Kashmir), India

Abstract

The present communication pertains to common weeds of rabi crops of tehsil Sunderbani, of the district Rajouri (Jammu & Kashmir), India. From the study area the 37 weed species belonging to 01 monocot and 16 dicot families are reported. Among dicot families the maximum dominance was shown by Asteraceae and Fabaceae, each represented by eight species. The only reported monocot family was Poaceae which included four weed species. The common weeds of rabi crops were *Avena fatua*, *Anagallis arvensis*, *Chenopodium album*, *Crisium arvense*, *Fumaria perviflora*, *Lathyrus aphaca*, *Melilotus indica*, *Parthenium hysterophorus*, *Phalaris minor*, *Rumex dentatus*, *Vicia hirsute* and *Vicia sativa* etc.

Keywords: Broadleaf weeds, grassy weeds, Rabi crops, identification.

Introduction

Weeds are unwanted plants that grow in association with agriculture crops and bring about significant decline in yield through their competition with crop plants for sunlight, space, nutrients etc. (Dangwal et al. 2010). However, some weeds are also allelopathic in nature (Oudhia and Tripathi, 1997 ; 1998). While Holm et al., (1977 ; 1979) estimated that about 8000 weed species growing in world, only 250 are of particular importance to agricultural crops. In view of significant yield decline by weeds in different crops, numerous studies have been carried out on various aspects of weed biology and control. The most fundamental of these studies is to document the composition of weeds that grow with crops. It is in this context Shailey and Gaur 1993 studies the phytosociological association of crops and weeds of pauri district of Uttarakhand, India and recorded 180 weed species belonging to 50 angiosperm families. The dominant dicot families were Amaranthaceae, Apiaceae, Asteraceae and Brassicaceae, in their studies. Among monocot families Commelinaceae and Poaceae were dominant. Gupta et al., (2008) studies the dynamics of cereal crop weeds of doon valley with special reference to rice, maize and wheat fields.

They reported 151 weed species belonging to 118 genera and 31 families; 57 weeds were reported from rice, 77 from maize and 71 from wheat fields. Kaul (1986) studied the weed flora of Kashmir valley and reported 401 weed species belonging to 251 genera and 56 angiosperm families. Singh et al., (2007) studied the phytosociological association of weeds in winter crops of Kashmir valley at varying altitudes from 1500 to 2000 m asl. They reported maximum IVI of *Panicum* in brown mustard throughout Kashmir valley. They found maximum IVI of *Ranunculus* spp. in wheat field of high altitudes and *Capsella bursa-pastoris* in wheat fields above 1600m asl.

The present study area i.e. tehsil Sunderbani, District Rajouri, India is located at an elevation of 800 m and situated at an latitude of 33^o.048N' and longitude of 74^o.49N'. The boundary of tehsil Sunderbani is surrounded on the eastern side with Nowshera and Reasi blocks and in the northern side with Rajouri. Its southern and western boundaries are bounded with Nowshera & Lam Darhal. Wheat is the major rabi crop of Tehsil Sunderbani but along with wheat, onion and mustard are also grown on small scale. The economy of this Tehsil revolves around production of its cash crops but the per hectare yield of crops in this Tehsil is less as compared to other Tehsil Sunderbani of district Rajouri, due to many factors out of which the problem of weeds is also prominent. The management of weeds involves costs therefore, reduction in net returns.



Sajjad Ahmed Mir

Research Scholar,
Deptt. of Zoology
Bharath University
Adyar, Chennai
Tamil Nadu,

Moreover, weeds in the study area make harvesting and threshing of crops costly, laborious and reduce the value of produce.

Objective of the Study

The present study was undertaken to find out common weeds of rabi crops in tehsil Sunderbani, Block Sunderbani of district Rajouri (Jammu & Kashmir), India.

Materials and Methods

Extensive and intensive field surveys were conducted during different months of rabi (winter) season (2009 – 10) in important villages of tehsil Sunderbani i.e. Malchola, Goder, Thandapani, Baja Bai, Rahsiote, Patrara, Chaniparat, Siote, Bal, Bakher, Taliri, Tali Gala and Mela etc. Three sites were selected in each village. Weeds were collected from all the sites of the study area. During this period interviews of farmers and Agriculturists were conducted to collect information about the seasonal weed plants and their vernacular names.

Result and Discussion

In all 37 weed species belonging to 01 monocot and 16 dicot families (Table -1) were collected from five villages of tehsil Sunderbani, District Rajouri (Jammu & Kashmir), India. The predominance was shown by Asteraceae and Fabaceae included five weed species and Poaceae was represented by four weed species. Amaranthaceae, Polygonaceae and Ranunculaceae were represented by three weed species each. Brassicaceae, Caryophyllaceae, Malvaceae and Solanaceae included two weed species each. The remaining families i.e. Asclepiadaceae, Cannabiaceae, Caesalpiniaceae, Convolvulaceae, Fumariaceae, Oxalidaceae, Primulaceae, Rosaceae and Rubiaceae were represented by one weed species each. The relative percentage of the different families exhibited 15.3% each for family Asteraceae and Fabaceae (Fig. 1). It is astonishing to note that grasses existed only to the extent of 7.6% among the weed flora of the target site. Out of 37 weed species reported from the study area, weeds like *Avena fatua*, *Anagallis arvensis*, *Chenopodium album*, *Cirsium arvense*, *Fumaria parviflora*, *Lathyrus aphaca*, *Melilotus indica*, *Parthenium hysterophorus*, *Phalaris minor*, *Rumex dentatus*, *Vicia hirsute* and *vicia sativa*

were common weeds of rabi crops in the study area. The weeds like *Euphorbia dracunculoides*, *Lolium temulentum*, *Polygonum barbatum*, *Polygonum persicaria* and *Ranunculus scleratus* were reported particularly from irrigated fields.

Some weeds reported from the study area, such as *Achyranthus aspera*, *Calotropis procera*, *Cannabis sativa*, *Chenopodium album* and *Cynodon dactylon* etc are of medicinal importance. The weeds like *Amaranthus viridis*, *Chenopodium album*, *C.vulgare*, *Coronopus didymus*, *Lathyrus aphaca*, *Vicia hirsute* and *V.sativa* are used in cooking recipes by Gujjar tribes of the study area. Fruit of *fragaria indica* are also edible. The present study may be helpful in identification of some common weeds of rabi crops. It may be helpful for taxonomists, agriculturists and scientists involved in the management of weeds.

The 01 monocot and 16 dicot families are arranged alphabetically with their botanical names, available vernacular names and flowering and fruiting season are mentioned (Table -1). These Findings are in a greater analogy with the previous work of Kaul (1986) who studied the weed flora of Kashmir valley and reported 401 weed species belonging to 251 genera and 56 angiosperm families. Since the present work was localized limited only to one tehsil, hence the flora is less diverse. Singh et al. (2007) studies the phyto-sociological association of weeds in winter crops of Kashmir valley at varying altitudes from 1500 to 2000 m asl. They reported maximum IVI of *Poa annua* in brown mustard throughout Kashmir valley. They found maximum IVI of *Ranunculus* spp. in wheat fields of high altitudes and *Capsella bursa-pastoris* in wheat could be attributed to the varying altitudes studied by those scholars. Hussain et al. (2004) studied the flora of tehsil Sunderbani, District Rajouri, and ten top most weeds in decreasing order of importance were *Mentha royleana* Benth., *Convolvulus arvensis* L., *Hordeum spontaneum* C. Koch. and *trifolium repens* L, etc. moreover, the recent studies of Hussain et al. (2009) also show a varying flora from the present findings due to the difference in agro-ecological conditions in the study areas. Our findings corroborated with the work of Swami and Gupta (1998).

Table-1. The 37 Weed Species Along with their Families, Available Vernacular Names and Flowering and Fruitinseasons.

S.No	Family	Botanical Name	Vernacularnae	Flowering & fruiting Season
1	Asteraceae	1. <i>Bidens pilosa</i> L. 2. <i>Cirsium arvensa</i> Syn. 3. <i>Conyza ambigua</i> Dc. 4. <i>Conyza bonariensis</i> L. 5. <i>Parthenium hysterophorus</i> L. 6. <i>Silybum marianum</i> L. 7. <i>Sonchus asper</i> L. 8. <i>Taraxacum officinale</i> weber.	Saryala Kandyari Booti Booti Congress booti kantili badi hand Hand	Sep. – Nov Jan. – March April – May Aug. – Sept. June – Sept.
2	Amaranthaceae	1. <i>Achyranthes aspera</i> L. 2. <i>Amaranthus spinosus</i> L. 3. <i>Amaranthus viridus</i> L.	Pooth kanda Chelari Chelari	Aug. – Dec. Sept.- Oct. Aug. – Nov.
3	Asclepiadaceae	1. <i>Calotropic procera</i> (Ait.) F.	Aak	April –July
4	Brassicaceae	1. <i>Capsella bursa-pastoris</i> Medik.	-----	Jan. – March

		2. <i>Coronopus didymus</i> (L.) Smith	-----	April. – Oct.
5	Caesalpiaceae	1. <i>Cassia occidentalis</i> L.	-----	Sept. – Nov.
6	Cannabiaceae	1. <i>Cannabis sativa</i> L.	Bhang	July – Sept.
7	Caryophyllaceae	1. <i>Silene conoidea</i> L. 2. <i>Stelleria media</i> L.	Doda ghash -----	March – April Feb. – March
8	Chenopodiaceae	1. <i>Chenopodium album</i> L. 2. <i>Chenopodium murale</i> L.	Bathua Laal bathua	April – June May – Oct.
9	Convolvulaceae	1. <i>convolvulus arvensis</i> L.	Bill	April – Sept.
10	Euphorbiaceae	1. <i>Euphorbia dracunculoides</i> Lamk. 2. <i>Euphorbia geniculata</i> Orteg. 3. <i>Euphorbia helioscopia</i> L. 4. <i>Euphorbia hirta</i> L. 5. <i>Euphorbia prostrate</i> Ait.	Doodle Badi doodal Doodal Choti doodi Doodi	Nov. – Jan. may – June May – July Sept. – Oct. July – Sept.
11	Fumariaceae	1. <i>Fumaria parviflora</i> Lamk.	Daniya ghash	Sept. - Nov
12	Malvaceae	1. <i>Malva Parviflora</i> L. 2. <i>Malvastrum coromen delianum</i> Syn.	Sonchal -----	March – April May – Aug.
13	Oxalidaceae	1. <i>Oxalis corniculata</i> L. Syn.	-----	Feb. – Nov.
14	Poaceae	1. <i>Avena fatua</i> L. 2. <i>Cynodon dactylon</i> (L.) Pers 3. <i>Lolium temulentum</i> L. 4. <i>Phalaris minor</i> Retz.	Gandial Ghash ----- Sitti	March – April April – July March – April March – April
15	Primulaceae	1. <i>Anagallis arvensis</i> L.	Krishna neel	Feb. – April
16	Rubiaceae	1. <i>Gallium aparine</i> L.	Chechra	Feb. – March
17	Rosaceae	1. <i>Fragaria indica</i> Andrews	Laal akhra	March – May

References

- Dangwal, L.R., A. Singh, T. Singh, A. Sharma and C. Sharma. 2010. effect of weeds on the yield of wheat crop in Tehsil Nowshera. J. American Sci. 6(10): 405 – 407.
- Gupta, A., S.P Joshi and R.K. manahas. 2008. multivariate analysis of diversity and composition of weeds communities of wheat fields in Doon valley. Ecological research Lab., Department of batony, D.A.V. (P.G). College, Dehradun (248007) Utrakhand India.
- Holm, L., J. Pancho, J. Herberger and D. Plucknett. 1979. A Geographical Atlas of World Weeds. John Wiley and sons, New York, 391 pp.
- Holm, L., D. plucknett, J. Herberger. 1977. The World's worst weeds: Distribution and Biology . University of Hawaii Press, Honolulu, 609 pp.
- Hussain, F., A. Murad and M.J Durrani. 2004. Weed communities in the wheat fields of Mastuj District Chitral, Pakistan. Pak. J. Weed Sci Res.10(3-4) : 101 -108.
- Hussain, F., S.M. Shah, Fazal-e-Hadi and Asadullah. 2009. Diversity and ecological characteristics of weeds of wheat field o University of Peshawar Botanical Garden at Azahkel, District Nowshera, Pakistan. pak. J. weed Sci. Res. 15 (4) : 283-294.
- Kaul, M.K 1986. weed flora of Kashmir valley. Journal of economics and taxonomic Botany. additional seriesscientific publishers, Jodhpur, India.
- Oudhia, P. and R.S. Tripathi. 1997. Allelopathic effects of *Parthenium hysterophorus* L. on kodo, Mustard and problematic weeds. First Intern. Conf. on *Parthenium* management. 11. UAS, Dhaward. India . 6-8 Oct. 1997,pp. 136 -139.
- Oudhia, P. and R.S. Tripathi. 1998. Allelopathic potential of *Datura stramonium* L. Crop Res. 16(1) , 37-40.
- shailey and R.D. Gaur. 1993. Phyto-sociological studies of cropsandweed association of Pauri district. Department of Botany. H.N.B Garhwal University . Utrakhand , India.
- Sharma , B.M and P. kachroo. 1983. Flora of Jammu and plants of neighborhood. Bishen Singh, Mahendrapal Singh, Dehradun, India.
- Singh, K.N., Ara, G.A. Wani, B. Hasan and B.A. Khanday. 2007. A phyto-sociological association of weeds in winter crops of Kasmir valley. Indian J. Weed Sci. 39(1-2):
- Swami, A. and B.K. Gupta. 1998. Flora of Udhampur. In B. Singh and M.Singh (eds.) Dehradun, India.