

A Study on Aflatoxin Contamination and Induced Biochemical Changes in *Syzygium Cumini*

Abstract

Mycoflora incidence and mycotoxin contamination in fruits/seeds of *Syzygium cumini* (L.) skeels collected from different localities of Uttarakhand (India) were investigated. A total of seven fungi namely *Aspergillus flavus*, *A. niger*, *A. repens*, *Fusarium avenaceum*, *F. semitatum*, *Rhizopus stolonifer* and *Trichothecium roseum* were recorded on fresh and stored samples of *Syzygium cumini*. *Aspergillus flavus* and *A. niger* was predominant. Nine percent of *Aspergillus flavus* isolates were toxigenic and produced aflatoxin B₁ in the up to 6.2 µg/ml in the liquid media. Results of natural occurrence of mycotoxins revealed that twenty-five percent of the stored samples was naturally contaminated with aflatoxins and the range of aflatoxin produced was 0.46-0.93 µg/g. Out of 17 positive samples, 14 samples exhibited presence of aflatoxin B₁, 2 samples aflatoxin B₁, B₂ and one sample exhibited presence of aflatoxin B₁, B₂ and G₁. A significant reduction in the level of alkaloid, starch and protein content of *S. cumini* seeds was recorded during fungal infection.

Keywords: *Syzygium Cumini*, Mycoflora, *Aspergillus Flavus*, Aflatoxins, Biochemical Changes

Introduction

Syzygium cumini (L.) skeels commonly known as Black plum or Jamun is one of the most important forestry species widely used as herbal medicine. It is an evergreen tropical tree of family Myrtaceae associated with many health and medicinal benefits. Various parts of *Syzygium cumini* are used for medicinal purpose such as bark, fruit, seed and leaves. While the bark and seeds are known for their hypoglycemic, anti-inflammatory, anti-diarrhoea and diuretic properties, leaves are used to treat diabetes, constipation etc. However, one of the best medicinal benefits of *S. cumini* seeds is its anti-diabetic properties. The seeds are reported to contain alkaloid, jambosine, and glycoside jambolin (antimellin), which halts the diastatic conversion of starch into sugar. Besides multiple medicinal properties, black plum is an important summer fruit having many vital nutrients like carbohydrates, protein, dietary fibers, fats etc. (Swami 2012).

Review of Literature

Mycotoxin contamination of various foodstuffs i.e. agricultural produce, forest produce and tree borne oil seeds has already been established (Hesseltine 1974; Bilgrami and Sinha 1984; Singh *et al.* 2001; Shukla and Singh 2006; Singh 2012). Medicinal plant materials have also been found prone to fungal infection and mycotoxin contamination and reports are available on presence of mycotoxin producing fungi and natural occurrence of mycotoxins in medicinal plants and traditional herbal medicines (Rani and Singh 1990; Roy and Chourasia 1990; Singh *et al.* 1999; Singh 2003). These plant materials get contaminated during harvesting, transportation, processing, storage or while still in the field. It has been established that fungal infection and mycotoxin contamination cause deterioration in the chemical constituents of the plant material under storage which may result in reduced efficacy and unsafe herbal preparations (Dutta and Roy, 1987). Gupta *et al.* (2016) also reported fungal contamination in powdered samples of leaves, stem bark and seeds/fruits of *S. cumini*.

The chemical composition and antioxidant activity of *S. cumini* have been studied in detail (Warrier *et al.* 1996; Benherlal and Arumughan 2007; Jadhav *et al.* 2009; Affity *et al.* 2011), but there is scant information available on mycotoxin contamination and induced biochemical changes in *S. cumini*.

Punam Kumari Singh
Regional Director,
IGNOU Regional Centre,
Shimla

Aim of The Study

To investigate aflatoxin contamination and induced biochemical changes in *S. cumini*.

Material and Methods

Samples collected from different forest areas and storage centres/ pharmaceutical industries of Uttarakhand (India) were screened for incidence of mycoflora, analyzed for the natural occurrence of aflatoxins and biochemical changes due to fungal contamination and mycotoxin production.

Mycoflora analysis

Mycoflora associated with the seed samples were examined using the blotter test and the agar plate method as recommended by International Seed Testing Association (ISTA, 1999). The developing fungal colonies were isolated, identified and maintained on potato dextrose agar (PDA) and Czapek's-dox agar media. Percentage incidence of fungi was determined on the basis of occurrence of a particular species in hundred seeds/samples.

Natural occurrence of Aflatoxins and toxigenic producing potential of *Aspergillus flavus*

Aflatoxin producing potentials of *Aspergillus flavus* isolates were tested in SMKY liquid medium (Diener and Davis, 1966). The constituents of the medium were, Yeast extract-7g; Sucrose- 200g; Magnesium Sulphate ($MgSO_4 \cdot 7H_2O$)-0.5g; Potassium Nitrate (KNO_3)-3g; Distilled water-1litre. Culture filtrate was extracted with chloroform and the chloroform extract was used for qualitative and quantitative detection of aflatoxins.

Samples were extracted chemically for the presence of aflatoxins by standard procedure (AOAC, 1984). Samples were powdered in a grinder and 50 g flour was blended with 250 ml methanol: water (60:40,v/v) for 2 minutes at high speed. The extract was filtered through Whatman No. 1 filter paper. 125 ml of this filtrate was then extracted with 30 ml saturated sodium chloride (NaCl) solution and 50 ml n- hexane in a 250 ml separating funnel for 2 minutes. The lower methanol layer was transferred to another separating funnel. Finally, the lower methanol layer was extracted with 40 ml chloroform. The chloroform extract was used for qualitative detection and quantitative estimation of aflatoxins. Qualitative and quantitative estimation of the mycotoxins were carried out using Thin Layer Chromatography (TLC). The spots of aflatoxins were identified by comparing with the spots of standards obtained from Sigma, USA.

To study the changes in alkaloid, protein and starch content of the *S. cumini* seeds, surface-sterilized seed samples were inoculated with spore suspensions (250-300 spores/ml) of *A. flavus* and incubated for 11 days at 96% r.h. Estimation of alkaloid, protein and starch content in the healthy and infested samples was done according to the standard recommended procedures (Harborne, 1998).

Results of mycoflora incidence, aflatoxin estimation and biochemical parameters was statistically analyzed using SPSS (1977) package. Standard error of means, t-value and critical difference were computed to draw appropriate inferences.

Results

A total of seven fungi namely *Aspergillus flavus*, *A. niger*, *A. repens*, *Fusarium avenaceum*, *F. semitectum*, *Rhizopus stolonifer* and *Trichothecium roseum* were recorded on fresh and samples of *Syzygium cumini*. While fresh fruits exhibited presence of only two fungi *A. niger* and *Fusarium avenaceum*, six fungi *A. flavus*, *A. niger*, *A. repens*, *F. semitectum*, *R. stolonifer* and *T. roseum* were recorded on stored samples. Occurrence of *A. niger* was the most common on all samples followed by *A. flavus*. While *A. candidus* was recorded only in monsoon season, *T. roseum* was recorded in summer and monsoon season only. Source and seasonal interaction was significant for *A. flavus*, *F. semitectum*, *R. stolonifer* and *T. roseum* and non-significant for *A. niger* and *A. candidus* (Table-1).

Aflatoxins were elaborated by *A. flavus* isolated from samples of *Syzygium cumini*, 3 isolates produced aflatoxin B₁ out of 32 screened and the amount of aflatoxin B₁ produced by the toxigenic isolates of *A. flavus* was in the range of 0.3-6.2 µg/ml. Results of natural occurrence of mycotoxins revealed that aflatoxins were not detected in fresh samples of *Syzygium cumini* while 25% of the stored samples were naturally contaminated with aflatoxins. Out of 17 positive samples, 14 samples exhibited presence of Aflatoxin B₁, Aflatoxin B₁,B₂ in 2 samples and one sample exhibited presence of Aflatoxin B₁,B₂ and G₁ and the range of aflatoxin production was in the range of 0.46-0.93 µg/g (Table-2).

Changes in the level of starch, protein, and total alkaloid during infestation of toxigenic strain of *Aspergillus flavus* have been carried out. *A. flavus* caused considerable changes in the starch, protein, and total alkaloid content during their infestation (Table-3). The amount of starch content in healthy seeds/ fruits was 11.15g/100g, however, the amount of starch in infested substrates was 9.61g/100g. Protein level was also depleted due to the infestation with *A. flavus*. The amount of protein content in healthy fruits/seeds was 1.03 g/100g, whereas in the infested sample, the amount of protein was 0.94 g/100g. There was an indication of inhibition in the total alkaloid content due to infestation with toxigenic strain of *A. flavus*. The alkaloid content decreased from 0.047% in control samples to 0.044 g/100g in the *A. flavus* infested samples. Statistical analysis of the results showed decline in the level of starch, protein, and total alkaloid significant at 5% level of significance.

Discussion

The results of the present investigation indicate that *S.cumini* seeds are subject to fungal contamination including that of *Aspergillus flavus*, a known mycotoxin producing fungi. Seasonal variation directly had an impact of percentage incidence of fungal flora. These medicinal plant produce get contaminated by various fungi during the process of harvesting, storage and further distribution causing spoilage and reducing the efficacy of the produce. Occurrence and incidence of mycoflora on a certain commodity is directly governed by the nature of the substrates, methods of storage and prevailing

environmental conditions. Earlier reports also indicate a varied pattern of fungal incidence and mycotoxin contamination in important medicinal species including *Aegle marmelos*, *Azadirachta indica*, *Emblica officinalis*, *Terminalis bellerica* and *T.chebula* (Aziz et al. 1988; Singh et al.1999; Chourasia 1995). Gupta et al.(2016) also reported fungal contamination in powdered samples of leaves, stem bark and seeds/fruits of *S. cumini*.

Screening of *A. flavus* isolates showed presence of Aflatoxin B₁ in higher concentration, besides, natural occurrence of aflatoxins is also quite high. The natural contamination of mycotoxins in these samples correspond to the incidence of toxigenic fungi associated with them and their respective potentiality to produce aflatoxins in the synthetic media. *Aspergillus flavus* group of fungi cause deterioration in the nutritive quality and medicinal property of the plant produce by producing aflatoxins and by changing some of the chemical constituents. Marked depletion in the alkaloid, starch and protein was recorded due to infestation with *A. flavus*. Aflatoxins and aflatoxin producing fungi have been reported to inhibit the synthesis of protein during plant metabolism (Truelove et al.1970; Dashek and Llewellyn 1974). Aflatoxin producing fungi have been known to degrade the protein level of some common dry fruits and fleshy fruits (Bilgrami et al.1983; Abdel-Hafez and Sabah 1993). Singh (2008) reported deterioration in nutritive quality of walnut kernels due to fungal infection and mycotoxin contamination. There is an indication of decline in the level of total alkaloid in samples infested with toxigenic strain of *A. flavus*. Inhibition in the level of total alkaloid might be due to their utilization or degradation into simpler forms. Dutta (1988) worked on deterioration in total alkaloid content of *Strychnos vomica* by some fungi and reported that *A. flavus* and *P. citrinum* significantly inhibited the level of alkaloid content.

Conclusion

Presence of mycotoxin producing fungi and natural occurrence of aflatoxins in the fruits/seed samples of *S. cumini* beyond tolerance level is a matter of great concern as it forms basic constituent of a large number of ayurvedic preparations. High incidence of naturally occurring aflatoxins in medicinal plant produce is a great threat to the quality and safety of the medicines. As a substantial part of the demand of herbal based pharmaceutical industries is met through wild collections from the forests, there is an urgent need to develop a mechanism to screen the samples for presence of toxigenic fungi and analyze for their level of mycotoxins before being consumed or marketed for use in the pharmaceutical industries.

Acknowledgement

The author is thankful to the Director, Forest Research Institute, Dehradun (Uttarakhand), India for the necessary laboratory facilities to carry out the research work. Financial assistance provided by Council of Scientific and Industrial Research (CSIR), New Delhi is gratefully acknowledged.

References

1. AOAC. 1984. *Official Methods of Analysis*. Association of Official Analytical Chemists, VA, USA, pp. 477-500.
2. Abdel-Hafez, A.I.I. and Sabah, M. Saber. 1993. *Mycoflora and mycotoxin in hazelnut (Corylus avellana) and walnut (Juglans regia) seeds in Egypt*. *Zentralbl. Mikrobiol.* 148 (2) :137-147.
3. Affity, A.M.R., Sayed, A.F., Shalaby, E.A., El-Shemy, A.H. 2011. *Syzygium cumini (pomposia) active principles exhibit potent anticancer and antioxidant activities*. *African J. Pharmacy and Pharmacology*, 5(7):948-956.
4. Aziz, N.H.; Youssef, Y.A.; El-Fouly, M.Z.; Moussa, L.A. 1998. *Contamination of some common medicinal plant samples and spices by fungi and their mycotoxins*. *Bot. Bull. Acad. Sinica.*, 39(4):279-285.
5. Benherhal, PS, Arumughan, C. 2007. *Chemical composition and and invitro antioxidant studies on Syzygium cumini fruits*. *J. Sci. Food Agric.*, 87:2560-2569.
6. Bilgrami, K.S., Sinha, KK and Singh, A. 1983. *Chemical changes in dry fruits during aflatoxin elaboration by Aspergillus flavus Link ex Fries*. *Current Science.*, 52 (10): 960-964.
7. Bilgrami K S and Sinha, KK. 1984 *Mycotoxin contamination of food and its control*. *Indian Rev Life Sci* 4:19 - 36.
8. Chourasia, H.K. 1995. *Mycobiota and mycotoxins in herbal drugs of Indian pharmaceutical industries*. *Mycol. Res.* 99 (6): 697- 703.
9. Dashek, W. V. and Llewellyn, G.C. 1974. *The influence of the carcinogen aflatoxin B1 on the metabolism of the germinating lilly pollen. In Fertilization in higher plants (Ed. H. F. Linskens). North- Holland Pub., Amsterdam: 351-360.*
10. Diener, U.L. and Davis, N.D. 1966. *Aflatoxin production by isolates of Aspergillus flavus*. *Phytopath.*, 56 (12) :1390-1393.
11. Dutta, G.R. and Roy, A.K. 1987. *Mycoflora associated with Strychnos seeds and deterioration of their active principles under storage*. *Indian Phytopathol* 40:520–524.
12. Gupta, M. Sharma, Sushil and Bhadauria, R. 2016. *Fungal contamination of powdered samples of different plant parts of two important medicinal plants: Momordica charantia and Syzygium cumini*. *Int. J. Pharm. Sci. Rev. Res.*, 38(1):179-184 .
13. Harborne, J.B. 1998. *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis (Chapman and Hall International Edition, New York) 3rd edition*.
14. Hesseltine CW. 1974. *Natural occurrence of mycotoxins in cereals*. *Mycol Appl* 53:141-153.
15. *International Seed Testing Association*. 1999. *Seed Sci Technol* 27 (Suppl).
16. Jadhav, VM., Kamble, S.S., Kadam, V.J. 2009. *Herbal medicine: Syzygium cumini :A Review*. *Journal of Pharmacy Research*, 2(8):1212-1219.
17. Rani, N. and Singh, S. 1990. *Aflatoxin contamination of some umbelliferous spices of human use*. *Int. Symp. aWorkshop on Food Cont.*

Mycotoxins and Phytotoxins, Abst. Book, pp 79-80.

18. Roy, AK and Chourasia, H.K. 1990. Mycoflora, mycotoxin producibility and mycotoxins in traditional herbal drugs from India. *J. Gen. Appl. Microbiol.* 36: 295-302.

19. Shukla, A.N. and Singh, Punam K. 2006. Fungal infestation and aflatoxin contamination in in edible oilseeds of forest origin. *J. Mycol & Plant Pathology* 36(1):96-100.

20. Singh, Punam Kumari, Khan, S.N. and Tandon, K.1999. Seed mycoflora of *Azadirachta indica* and its control. *Van Vigyan*, 37: 18-20.

21. Singh, PK, Khan SN, Harsh NSK and Pandey,R. 2001. Incidence of mycoflora and mycotoxins in some edible fruits and seeds of forest origin. *Mycotoxin Research* 17(2) 46-58.

22. Singh, Punam Kumari 2003. *Mycotoxin elaboration in Triphala and its constituents*. *Ind. Journal of Phytopathology*, 56(4), 380-383.

23. Singh, Punam Kumari and Shukla, A.N. 2008. Survey of mycoflora counts, aflatoxin production and induced biochemical changes in walnut kernels. *J. Stored Product Research*, 44: 169-172.

24. Singh, Punam K. 2012. Assessment of mycotoxins in Tree Borne Oil Seeds, *J. Food Research*, Vol.1 (3):92-101.

25. Swami, S.B., J.Thakor, N.S., Patil, M., and Haldankar, Parag M. 2012. Jamun (*Syzygium cumini* (L.): A Review of Its Food and Medicinal Uses. *Food and Nutrition Sciences*, 2012, 3:1100-1117.

26. Truelove,B., Davis,D.E.and Thompson,O.C.1970. Effects of aflatoxin B1 on protein synthesis by cucumber cotyledon discs. *Can. J. Bot.*, 48: 485.

27. Warriar,PK., Nambiar,VPK., Ramankutty, C. 1996. *Indian Medicinal Plants*. Orient Longman Ltd. Hyderabad India).5:225-228.

29. Wikipedia: https://en.wikipedia.org/wiki/Syzygium_cumini

Table-1 Mycoflora Incidence in Stored Samples of *S. Cumini*

Mycoflora	Mean Value of Mycoflora Incidence in Different Season			
	Season	Mean	Effect	CD _(0.05)
<i>A. flavus</i>	Summer	43.75	Season	1.17
	Monsoon	47.5	Source	1.36
	Winter	17.15	Season x Source	2.35
<i>A. candidus</i>	Summer	0.00	Season	0.76
	Monsoon	18.6	Source	NS
	Winter	0.00	Season x Source	NS
<i>A. niger</i>	Summer	46.95	Season	1.29
	Monsoon	47.95	Source	1.50
	Winter	15.25	Season x Source	NS
<i>F. semitectum</i>	Summer	11.25	Season	1.07
	Monsoon	11.15	Source	1.24
	Winter	3.15	Season x Source	2.15
<i>R. stolonifer</i>	Summer	14.55	Season	1.41
	Monsoon	12.1	Source	1.63
	Winter	6.90	Season x Source	2.82
<i>T. roseum</i>	Summer	2.35	Season	0.71
	Monsoon	8.15	Source	0.82
	Winter	0.00	Season x Source	1.41

Table-2 Natural occurrence of aflatoxins in *S. cumini*

Total No. of samples		Percentage contamination	Aflatoxins detected	No. of samples	Conc. of Aflatoxin B ₁ (µg/g)
Screened	Contaminated				
68	17	25	Aflatoxin B ₁	14	0.46- 0.93
			Aflatoxin B ₁ ,B ₂	2	
			Aflatoxin B ₁ ,B ₂ ,G ₁	1	

Table-3: Changes in Starch, Protein and Alkaloid Content Due to Fungal Infection

Biochemical parameters		g/100g (Mean±S.E.)	t-value	Significance CD _(0.05)
Starch	Control	11.15±0.0091	7.85	0.004
	Infested	9.612±0.1903		
Protein	Control	1.032±0.0811	8.73	0.001
	Infested	0.938±0.0759		
Alkaloid	Control	0.0472±0.000	21.36	0.022
	Infested	0.0435±0.000		